

On the admissibility of topological vector spaces

O. HADŽIĆ

1. Introduction. Let X be a Hausdorff topological vector space. A subset A of X is called *admissible* [7] if for every compact subset $K \subset A$ and for every neighbourhood U of zero in X there is some continuous mapping $h: K \rightarrow A$ such that

- (i) $\dim(\text{span } h(K)) < \infty$,
- (ii) $x - hx \in U$, for all $x \in K$.

S. HAHN and K. F. PÖTTER [3] proved fixed point theorems for admissible subsets of Hausdorff topological vector spaces. NAGUMO proved that all convex subsets of a locally convex space are admissible [9] and the admissibility of many non-convex topological vector spaces has been proved by KLEE [6], RIEDRICH [14], [15], ICHII [4], PALLASCHKE [12] and KRAUTHAUSEN [7].

But the following questions remained open:

- a) Which Hausdorff topological vector spaces are admissible?
 - b) Which convex subsets are admissible?
 - c) For which compact subsets K of a Hausdorff topological vector space X is the following valid:
- (*) If U is an arbitrary neighbourhood of zero in X then there is a finite dimensional continuous mapping $h: K \rightarrow \text{co } K$ such that $x - hx \in U$ for all $x \in K$.

Recently, MATUSOV [8] proved that every compact convex subset of a Hausdorff topological vector space has the fixed point property using an idea of SARIMSAKOV [10] and a result of KASAHARA [5].

Now we give Kasahara's definition of paranormed spaces [5].

A linear mapping Φ of a topological semifield E into another F is said to be *positive* if $\Phi(x) \geq 0$ in F for every $x \in E$ with $x \geq 0$. Let $\| \cdot \|$ be a mapping of a

linear space X into a topological semifield E and let Φ be a continuous positive linear mapping of E into itself. The triple $(X, \|\cdot\|, \Phi)$ is called a *paranormed space over E* and $\|\cdot\|$ a Φ -*paranorm* on X over E if the following conditions are satisfied:

- (P1) $\|x\| \geq 0$, for every $x \in X$;
 (P2) $\|\lambda x\| = |\lambda| \cdot \|x\|$ for every real λ and every $x \in X$;
 (P3) $\|x + y\| \leq \Phi(\|x\| + \|y\|)$ for every $x, y \in X$.

A set $K, K \subset X$ where X is a topological vector space, is said to be of type Φ iff $(X, \|\cdot\|, \Phi)$ is a paranormed space and for every $n \in \mathbb{N}$, every $x_1, x_2, \dots, x_n \in K - K$ and every $\lambda_i, 0 \leq \lambda_i \leq 1$ ($i = 1, 2, \dots, n$) such that $\lambda_1 + \lambda_2 + \dots + \lambda_n = 1$, we have $\left\| \sum_{i=1}^n \lambda_i x_i \right\| \leq \sum_{i=1}^n \lambda_i \Phi(\|x_i\|)$. If $K = X$, the space X is of type Φ .

In this paper we shall prove:

- a') Every Hausdorff topological vector space of type Φ is admissible.
 b') Every convex subset of type Φ of a Hausdorff topological vector space is admissible.
 c') For every compact subset K of type Φ of a Hausdorff topological vector space property (*) is valid.

As a Corollary we shall obtain an extension of Matusov's fixed point theorem.

2. The main result. We use the following theorem from KASAHARA's paper [5].

Let (X, τ) be a topological linear space. Then there exists a paranormed space $(X, \|\cdot\|, \Phi)$ over a Tihonov semifield E such that:

- (1) *For every neighbourhood U of $0 \in X$ there are an $\varepsilon > 0$ and an indecomposable idempotent $\varrho \in E$ such that*

$$\{x \in X: \|x\| \cdot \varrho \leq \varepsilon \varrho\} \subset U.$$

- (2) *For every neighbourhood U of $0 \in E$ the set*

$$\{x \in X: \|x\| \in U\}$$

is a neighbourhood of $0 \in X$.

The Tihonov semifield E from the above Theorem is R_{Δ} , the set of all mappings from Δ into R where Δ is a set of paranorms generating the topology of X and satisfying the condition that for each $p \in \Delta$ there are $\alpha > 0$ and $q \in \Delta$ such that

$$p(x+y) \leq \alpha(q(x)+q(y)), \text{ for all } x, y \in X.$$

Now we are ready to formulate our main theorem.

Theorem. *For every compact subset K of type Φ of a topological vector space X and for every neighbourhood U of zero in X there exists a finite dimensional continuous mapping $h: K \rightarrow \text{co } K$ such that $x - hx \in U$ for all $x \in K$.*

Proof. Let U be an arbitrary neighbourhood of zero in X and let $\mu = \{t_1, t_2, \dots, t_n\} \subset \Delta$ and $\varepsilon > 0$ such that

$$\|x - y\| \in U_{\mu, \varepsilon} \Rightarrow x - y \in U,$$

where

$$U_{\mu, \varepsilon} = \{u: u \in R_\Delta, u(t_j) < \varepsilon, j = 1, 2, \dots, n\}.$$

Further, since the mapping $\Phi: R_\Delta \rightarrow R_\Delta$ is a continuous linear mapping there exists a neighbourhood $V_1(\mu, \varepsilon)$ of zero in R_Δ such that

$$\|x - y\| \in V_1(\mu, \varepsilon) \Rightarrow \Phi(\|x - y\|) \in U_{\mu, \varepsilon}.$$

Suppose now that $V_2(\mu, \varepsilon)$ is a circled neighbourhood of zero in X such that

$$x - y \in V_2(\mu, \varepsilon) \Rightarrow \|x - y\| \in V_1(\mu, \varepsilon).$$

Since X is a Hausdorff topological vector space it is also a Hausdorff uniform space and let d be a pseudometric on X and $\delta > 0$ such that

$$d(x, y) < \delta \Rightarrow x - y \in V_2(\mu, \varepsilon).$$

We shall use the notation

$$V_x(d, \delta) = \{y: y \in X, d(x, y) < \delta\} \quad (\delta > 0).$$

Since the set K is compact there exists a finite set $\{x_1, x_2, \dots, x_m\} \subset K$ such that for every $x \in K$ there exists $i \in \{1, 2, \dots, m\}$ such that

$$x \in V_{x_i}(d, \delta).$$

So if we define the functions $f_i: K \rightarrow R^+$ ($i = 1, 2, \dots, m$) so that

$$f_i(x) = \max \{0, \delta - d(x, x_i)\}$$

for every $x \in K$ and $i \in \{1, 2, \dots, m\}$ it follows that

$$f_i(x) \neq 0 \Leftrightarrow d(x, x_i) < \delta.$$

Since for every $x \in K$ there exists at least one $i \in \{1, 2, \dots, m\}$ such that $f_i(x) \neq 0$ we conclude that for every $x \in K$,

$$s(x) = \sum_{i=1}^m f_i(x) \neq 0$$

and that all mappings f_i ($i=1, 2, \dots, m$) are continuous since the mapping $x \mapsto d(x, x_i)$ is continuous for every $i \in \{1, 2, \dots, m\}$. Now, let

$$h(x) = \frac{1}{s(x)} \sum_{i=1}^m f_i(x) x_i \quad \text{for all } x \in K.$$

Then $h(K) \subset \text{co } K$ and h is a continuous mapping from K into a finite dimensional subspace of X . Further we have

$$\begin{aligned} \|hx - x\| &= \left\| \frac{1}{s(x)} \sum_{i=1}^m f_i(x) x_i - \frac{1}{s(x)} \sum_{i=1}^m f_i(x) x \right\| = \\ &= \left\| \frac{1}{s(x)} f_i(x) (x - x_i) \right\| \leq \frac{1}{s(x)} \sum_{i=1}^m f_i(x) \Phi(\|x - x_i\|). \end{aligned}$$

Since $f_i(x) \neq 0 \Leftrightarrow d(x, x_i) < \delta$ it follows that for every $x \in K$ such that $f_i(x) \neq 0$ we have that

$$\Phi(\|x - x_i\|) \in U_{\mu, \varepsilon}$$

and so

$$\|hx - x\|(t) \leq \frac{1}{s(x)} \sum_{i=1}^m f_i(x) \Phi(\|x - x_i\|)(t) < \frac{1}{s(x)} \sum_{i=1}^m f_i(x) \varepsilon = \varepsilon \quad \text{for every } t \in \mu.$$

So we have $\|hx - x\| \in U_{\mu, \varepsilon}$, which implies $hx - x \in U$ and the proof is complete.

Corollary 1. *Every convex subset A of type Φ of a Hausdorff topological vector space is admissible.*

Proof. If K is a compact subset of A and U is an arbitrary neighbourhood of zero, the Theorem implies the existence of a finite dimensional continuous mapping $h: K \rightarrow \text{co } K$ with the following property:

$$x - hx \in U \quad \text{for all } x \in K.$$

Since A is convex it follows that $\text{co } K \subset A$ and so A is admissible.

Corollary 2. *Every Hausdorff topological vector space of type Φ is admissible.*

Corollary 3. *Let A be a closed and convex subset of type Φ of a Hausdorff topological vector space E and $h: A \rightarrow A$ be a continuous mapping such that $\overline{h(A)}$ is compact. Then there exists at least one fixed point of the mapping h .*

Proof. Since A is admissible we can apply a fixed point theorem from [3] and so the set of fixed points of the mapping h is nonempty.

References

- [1] O. HADŽIĆ, *The Foundation of the Fixed Point Theory*, Institut za matematiku (Novi Sad, 1978) 320 pp.
- [2] S. HAHN, A remark on a fixed point theorem for condensing setvalued mappings, *Informationen*, Technische Universität Dresden, 07—5—77.
- [3] S. HAHN—F. K. PÖTTER, Über Fixpunkte kompakter Abbildungen in topologischen Vektorräumen, *Stud. Math.*, **50** (1974), 1—16.
- [4] J. ISHII, On the admissibility of function space, *J. Fac. Sci. Hokkaido Univ.*, Ser. I, **19** (1965), 49—55.
- [5] S. KASAHARA, On formulations of topological linear spaces by topological semifield, *Math. Japonicae*, **19** (1974), 121—134.
- [6] V. KLEE, Leray—Schauder theory without local convexity, *Math. Ann.*, **141** (1960), 286—296.
- [7] C. KRAUTHAUSEN, On the theorems of Dugundjy and Schauder for certain nonconvex spaces, *Math. Balk.*, **4** (1974), 365—369.
- [8] V. E. MATUSOV, Obobščenie teoremy o nepodvižnoj točke Tihonova, *Doklady A. N. Uz SSR*, No 2 (1978), 12—14.
- [9] M. NAGUMO, Degree of mapping in convex linear topological spaces, *A. J. Math.*, **73** (1951), 497—511.
- [10] T. A. SARIMSAKOV, Novoe dokazatel'stvo teoremy Tihonova, *Uspehi M. N.*, **20** (124) (1965).
- [11] D. R. SMART, *Fixed point theorems*, Cambridge University Press (1974).
- [12] D. PALLASCHKE, Verallgemeinte Basen für topologische lineare Räume, *Schriften der Ges. Math. und Datenverarb.*, Bonn, **8** (1969).
- [13] J. REINERMANN—V. STALLBOHM, Fixed point theorems for compact and nonexpansive mappings on starshaped domains, *Comment. Math. Univ. Carolinae*, **15** (1974), 775—778.
- [14] T. RIEDRICH, Die Räume $L^p(0, 1)$ ($0 < p < 1$) sind zulässig, *Wiss. Z. Techn. Univ. Dresden*, **12** (1963), 1149—1152.
- [15] T. RIEDRICH, Der Raum $S(0, 1)$ ist zulässig, *Wiss. Z. Tech. Univ. Dresden*, **13** (1964), 1—6

DEPARTMENT OF MATHEMATICS
 FACULTY OF SCIENCE
 UNIVERSITY OF NOVI SAD
 ULICA DR ILIJE DJURIČIĆA 4
 21000 NOVI SAD
 YUGOSLAVIA